



MEDICATIONS TO AVOID PRIOR TO SURGERY

All patients that are scheduled for surgery should discontinue taking any and all products that contain aspirin and Vitamin E for two (2) weeks prior to your procedure.. Aspirin and Vitamin E may cause bleeding and/or clotting problems during and after surgery. **If your doctor has prescribed any of these medications, please contact us for instructions.**

The following is a list of medications that should be avoided:

Advil	Disalcid	Percodan
Alcohol (24 hours prior)	Doan's	Percodan-Demi
Aleve	Dolia	Persantine
Alka-Seltzer	Dolobid	Persistin
Anacin	Dristan	Quagesic
Anaprox	Duardyne	Relafen
Anaproxon	Duragesic	Robaxisal
Ansaid	Easprin	Rufen
APC	Ecotrin	Sine-Aid
Arthritis Pain Formula	Empirin	Sine-Off
ASA	Emprazol	Soma with Codeine
Ascriptin	Equagesic	Soma Compound
Ascodeen-30	Excedrin	Stanback Formula
Aspercream	Feldene	Max Powder
Aspergum	Fiorinal	Stendin
Aspirin	Fluriprophen Sodium	Stero-Darvon with ASA
Bayer Aspirin	Four Way Cold	Supac
BC Cold Powder	Goody's Extra Strength	SX-65 compound
BC Powders	Ibuprofen	Synalgos
Brufen	Indocin	Synalgos DC
Buff-A-Corn	Indomethacin	Talwin
Buffadyne	Magsal	Tolectin
Bufferin	Measurin	Toradol
Buffex	Meclomen	Trandate
Butabital	Medipren	Trental
Cama Arthritis Pain Reliever	Midol-200	Trialgesic
Cama-Inlay	Midol	Triamincin
Caprin	Midol PMS	Trilisate
Caspirin	Mobigesic	Ursinus Inlay
Cephalgesic	Monacet with Codeine	Vanquish
Cheracol	Momentum Muscle Back Form	Vivo Med
Children's Aspirin	Motrin	Vitamin E
Clinoril	Naprosyn	Voltaren
Conaterol	Norgesin	Wesprin
Congespirin	Norgesic Forte	Zactrin
Cope	Nuprin	Zorpin
Coricidin	Pabirin Buffered	Gingko
Coumadin	Panadynes	Rogaine
Darvon with ASA	Panalgesic	Pepto-Bismol
Darvon Compound	Daypro	Merida
Ginsing	Plavix	Effedra